

Of maize and men, or peas and people: case histories to justify plants and other model systems

David Baulcombe

One of the byproducts of molecular biology has been support for the 'model system' concept. All living organisms are based on the same genetic code, they have similar subcellular structures and they use homologous metabolic pathways. So, mechanisms can be investigated using organisms other than those in which the knowledge will be exploited for practical benefit. Model systems are particularly useful in the early discovery phase of a scientific endeavor, and recent progress in biomedical science has fully vindicated their use. Jacques Monod, for example, famously justified his work on a bacterial model system by stating that "what is true for *Escherichia coli* is also true for elephants." My fellow laureates, Victor Ambros and Gary Ruvkun, can defend the use of the worm *Caenorhabditis elegans* as a good model system and so I will focus on plants.

Probably the best example I could use is that of Barbara McClintock, who used corn in her Nobel Prize-winning work¹. Her identification of mobile genetic elements has been enormously influential in microbiology and immunology, and continues to bear on the field of epigenetics that I refer to again below. She could have adapted Monod's aphorism to refer to maize and men. Gregor Mendel would, of course, have referred to peas and people.

There are many other historical examples, in addition to McClintock's and Mendel's work, that could justify the use of plants as models. Robert Hooke used his newly developed microscope in 1664 to show that cork comprises structures that he referred to as cells ("Our microscope informs us that the substance of

cork is altogether filled with air, and that air is perfectly enclosed in little boxes or cells distinct from one another.")² (Fig. 1). Two hundred fifty years later, Beijerinck discovered a *contagium vivum fluidum* in extracts of diseased tobacco plants that he later referred to as a virus³.

In contemporary science, a green alga—*Chlamydomonas reinhardtii*—is a useful model in the analysis of kidney disease⁴. However, in this article, I refer to the contribution of plant biology to a family of mechanisms that I refer to as RNA silencing. This topic has been reviewed comprehensively elsewhere^{5,6}, so here I focus on personal experience and my view of future potential from this work.

The early history of RNA silencing in plants

The most important development in plant biology in the 1980s was the introduction of methods for transfer of DNA into the nuclear genome⁷. Transgenic techniques were developed as a routine research tool, and the new technology prompted the introduction of genetic manipulation for crop improvement. The first generation of genetically modified plants used bacterial genes for herbicide tolerance or insect resistance^{8,9}. There were also attempts from several laboratories, including my own, to develop virus resistance by transgenic expression of virus-derived genes in the host organism^{10–12}. We were testing a concept known as parasite-derived resistance in which the replication or spread of a disease agent could be reduced by transgenic expression of its genes in a host. A third category of these early transgenic crops exploited an antisense approach to block expression of an endogenous plant gene¹³. Crops were to be improved in this approach by specific suppression of an endogenous gene that limited agronomic performance. The idea was simple: a duplex RNA





Figure 1 Robert Hooke's micrograph of cells in a sample of cork. Downloaded from http://www.nlm.nih.gov/exhibition/historicalanatomies/Images/1200_pixels/hooke_t11.jpg.

structure would be formed between the mRNA of the target gene and a transgenic antisense RNA. This duplex structure would then either block translation of the mRNA or initiate its degradation by a nuclease specific for double-stranded RNA.

Many of the early transgenic plants had the desired traits, and they were the progenitors of transgenic crops that are now grown in large areas of different countries. However, there were anomalous phenotypes in some of the transgenic lines. For example, the bacterial and viral transgenes were not always expressed^{8,9}. In the 'antisense plants', the suppression effect

David Baulcombe is Royal Society Research Professor and Professor of Botany, Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, UK.
e-mail: dcb40@cam.ac.uk



Richard Jorgensen

Figure 2 Variegation of silencing in flowers of petunia. The target of silencing by a sense transgene is chalcone synthase. The silencing effect manifested as the absence of pigment is present in only part of the flower. Reprinted with permission of the American Society of Plant Biologists.

was sometimes variegated—it was present on some branches of the plant and not in others, or it was absent in localized parts of a flower¹⁴. There was also a problem with some of the sense RNA controls in these antisense experiments. There was a silencing effect even when the transgene was designed to generate sense RNA that would not form a duplex with the target mRNA^{15–17}. In some examples, there was variegation of this sense suppression as with the antisense plants (**Fig. 2**). Another strange process occurred when two transgenes carried the same promoter sequence: one of these transgenes could silence the other through a process operating at the transcriptional level¹⁸.

In retrospect, I can see that these anomalies challenged the existing paradigm of genetic regulation, but initially, I made the mistake of thinking that they were due to multiple distinct mechanisms and that they were artifacts of transgenes. However, I became more interested when we characterized transgenic plants manifesting parasite-derived resistance. The plants with the strongest viral resistance were those in which the transgene RNA was present at a low abundance, whereas plants expressing the same gene at a high level were fully susceptible¹⁹. It seemed that the transgene silencing in this example of ‘less is more’ could not be dismissed as a side effect or artifact.

A new model to explain some of the counterintuitive properties of the virus-resistant plants was attractive—at least to us—because it could account for the anomalous results with many other transgenic plants, although it could not explain the variegation or trans-inactivation effects. The model proposed that certain transgenes, irrespective of whether they were designed to be expressed in sense or antisense orientation, have a silencing property

that operates in a nucleotide sequence-specific manner. The silencing *in cis* would explain why the transgenes in certain lines were not abundantly expressed. Correspondingly, the silencing *in trans* would account for the ability of these transgenes to silence endogenous genes or viruses^{20,21}.

The hypothesis further proposed that the silencing mechanism operates at the RNA level because the targeted viruses had an RNA genome and did not have a DNA phase in their replication cycle. Originally, many of the silencing-related phenomena in transgenic plants were referred to as ‘gene silencing’. However, at a seminar in Switzerland, I was interrupted by Ingo Potrykus—the inventor of ‘golden rice’²²—who pointed out that I was not talking about ‘gene silencing’; it was ‘RNA silencing’. I agreed and have since promoted ‘RNA silencing’ as a generic term to cover a family of mechanisms involving silencing and RNA.

Support for the model came from a series of tests carried out by Jim English with genetically modified viruses²⁰, and the remaining challenges were to find out about the details of the mechanism and its biological role. We were also interested in finding out whether the variegated silencing of transgenes and transcriptional silencing of promoter sequences could be accommodated in the model.

A central mystery in RNA silencing was its ability to act in a nucleotide sequence-specific manner. The simplest explanation required an antisense RNA that would guide a silencing machinery to its target. In the transgenic experiments described above, this antisense RNA would be present irrespective of whether the transgene was in a sense or an antisense orientation. Andrew Hamilton, who had been interested in antisense RNA ever since his PhD work on transgenic tomatoes²³, joined the lab to carry out the search for this elusive molecular species. It is testimony to his persistence and talent as a scientist that he eventually found them²⁴. He tried several different methods, and his early successes used a sensitive but imprecise RNase protection method that generated rather ugly looking smears instead of the neat electropherograms that were eventually published in *Science* (**Fig. 3**). Our first estimate was that these antisense RNAs were 25 nucleotides long, but later, we refined this to 21–24 nucleotides²⁵. They are now known as small interfering RNAs (siRNAs)²⁶.

The initial convergence of our work with that of Ambros and Ruvkun was because the methods we used to detect siRNAs were the same as those used to detect small temporal RNAs from *lin-4* in worms^{27,28}. However, we did not immediately make the biological connection. In retrospect, I cannot say why—it seems obvi-

ous now. My failure to make the connection was probably because the mechanism in worms involved suppression of translation, whereas, in plants, we were dealing with a process that decreased stability of the targeted RNA.

Andrew Hamilton’s experiments further demonstrated that the production of small antisense RNA required transcription of the corresponding sense strand. This involvement of both RNA strands provided an obvious connection between RNA silencing in plants and RNA interference in worms as revealed by the famous Andrew Fire and Craig Mello 1998 paper in *Nature*²⁹. Once this connection was recognized, there was an opportunity for discoveries in plants to catalyze our understanding of animal systems and vice versa. It has been enormously satisfying to witness this exchange of information and the elucidation of variations on a molecular theme associated with different types of silencing mechanism in diverse organisms^{5,6}.

Much current RNA silencing research interest is focused on endogenous small RNAs. In plants, there are thousands of loci that produce these small RNAs^{30–32}. A small proportion of the endogenous small RNAs are similar to the microRNAs of worms that were first discovered as short temporal RNAs by Ruvkun and Ambros^{27,28}. Most microRNAs are negative regulators, although a positive regulatory microRNA has been recently described³³. The other small RNAs in plants are siRNAs, the double-stranded RNA precursors, which are produced in various ways, including the annealing of complementary RNA, foldback of long inverted repeats and the action of RNA-dependent RNA polymerases on a single-stranded RNA^{30,31}.

It could be said that, now that we have the understanding of multiple variations on silencing mechanisms, RNA silencing research with a biomedical target should focus exclusively on people or at least on vertebrate experimental systems. However, for this view to be valid, the discovery phase of RNA silencing research would have to be played out. I do not consider this to be the case just yet because there are at least two areas—virus resistance and epigenetics—in which there is much to be discovered and in which RNA silencing in plants has the potential to inform biomedical research.

RNA silencing and virus resistance

Genetic engineering of virus resistance was my introduction to RNA silencing, and there is a nice irony from the subsequent discoveries that plants are naturally protected from viruses by RNA silencing. Frank Ratcliff, Tamas Dalmay and Olivier Voinnet joined my group to investigate these aspects of RNA

silencing together with Andrew Hamilton. Their work, alongside that of the Covey, Carrington and Vance groups, revealed a picture in which virus-specific siRNAs are produced using a viral RNA template in infected plant cells^{34–40}. These siRNAs then target an RNA silencing effector protein to the viral RNA so that the replication cycle of the virus is blocked. Inevitably, in the arms race of defense and counter-defense between viruses and their hosts, plant viral genomes encode pathogenicity or virulence factors that are suppressors of silencing. Olivier's work, together with the work of those other groups, revealed that there are many different types of viral suppressor protein^{36–40}.

Associated with antiviral silencing, there is a mobile signal that mediates antiviral effects of silencing beyond the infected regions of the plant and that prevents or delays spread of the virus in the infected plant. The discovery of signaling^{41,42} was particularly satisfying because, in addition to virus resistance, it explained the variegation anomaly in the early transgenic antisense and sense RNA experiments: signal produced in one part of the plant spread unevenly so that the silencing effect was manifested only in the parts in which it was received.

We now understand that RNA silencing influences many aspects of the interactions between viruses and their plant hosts. It explains, for example, how viruses are excluded from the meristem of infected plants, why some plants recover from viral disease and why virus-infected plants may be protected from secondary infection. Many aspects of symptom production may also be due to suppression of endogenous genes by viral siRNA. Additionally, in a biomedical context and consistent with the model system justification for plant work, RNA silencing can target viral RNA in animals.

Invertebrates may use RNA silencing as part of natural antiviral defense, as in plants⁴³. In vertebrates, RNA silencing may not be part of the natural antiviral defense systems⁴⁴. However, artificial RNA silencing can be effectively targeted against viruses⁴⁵. Given the difficulty of developing small molecule therapeutic agents to target viruses, there is much interest in the possibility of delivering double-stranded RNA or siRNAs to protect against viral disease. A major challenge in such a task is to find ways to chemically modify the RNA so that it is stable until it is taken up by cells and targeted to the desired cell type. This challenge is therefore essentially pharmacological and probably outside the influence of plant biology. However, it may be possible, on the basis of an understanding of the plant viral systems, to optimize the

intracellular efficacy of antiviral siRNAs. Nature has already done the experiments in plants to find out which regions of viral genomes are the most effective targets of siRNAs and whether the antiviral potential is influenced by base composition or other chemical characteristics of the siRNAs. Given the structural and genetic similarities of various groups of plant and animal viruses, it is likely that the understanding of these plant systems will facilitate development of novel antiviral therapy in medicine.

RNA silencing and epigenetics

The final anomaly in the original transgenic experiments—promoter silencing at the transcriptional level—was finally resolved by the Matzke group and others. They demonstrated that double-stranded RNA and siRNAs have the potential to guide epigenetic modification of DNA or the associated chromatin proteins in plants and other organisms^{25,46,47}. The silencer transgene evidently produces siRNAs that can target the promoter of the silenced gene.

There is now a good understanding of this epigenetic RNA silencing mechanism⁴⁸, and we also know that many endogenous siRNAs are associated with epigenetic modification of the corresponding DNA^{30,49}. The siRNA associated with epigenetic modification is 24 nucleotides long. However, consistent with the past history of RNA silencing research, a mystery remains because these 24-nucleotide siRNAs do not have an obvious role. Mutant plants that do not produce these 24-nucleotide siRNAs are able to grow and develop at a normal rate⁵⁰. We can therefore rule out the possibility that these RNAs have a major role in normal growth and development. Many of these endogenous siRNAs correspond to transposons³⁰. However, transposons are not mobilized in RNA silencing mutants, and it seems unlikely that these RNAs have the obvious role in protection against damage caused by mobile genetic elements.

Resolution of this mystery may not have immediate relevance to biomedicine because the answer may involve mechanisms in evolution. One possibility is that epigenetic RNA silencing by one parent may silence essential genes in the opposite parental genome and thereby influence hybrid formation. A second possible role is prompted by Louise Jones's finding that RNA-directed epigenetic changes may persist for several generations even in the absence of the initiator RNA⁵¹. This finding illustrates how there are separate initiation and maintenance phases to the RNA-directed epigenetic mechanism. It also illustrates how endogenous siRNAs that are induced by environmental stimuli could induce heritable phenotypic changes. Such changes would not be associated with genetic mutation but nevertheless could be subject to




Figure 3 One of Andrew Hamilton's early RNase protection assays showing that small antisense RNA—the black smear corresponding to 22 nucleotides (nt)—was only present in plants that were silencing the target RNA. Production of the *GUS* reporter gene antisense RNA was suppressed if the sense transcription was blocked. One of Andrew Hamilton's early RNase protection assays showing that small antisense RNA was only present in plants that were silencing the target RNA. Production of the *GUS* reporter gene antisense RNA, detected by the anti-*GUS* RNA probe, was suppressed if the sense transcription was blocked. Far left, hybridization probe without RNase treatment. The two unmarked samples in the next two lanes were from plants without a *GUS* reporter gene; there is no evidence of any antisense RNA. The other three samples (white symbols) were from plants with a *GUS* transgene. The sample on the far right (blue symbol) had an untranscribed *GUS* transgene and there was also no antisense RNA. However, the two samples from a plant with a transcribed *GUS* transgene (red symbols) that was silenced at the RNA level generated a smeared signal centered on the 22-nucleotide length that was the first evidence for siRNAs.

selection in certain environments.

If this RNA-directed epigenetic modification occurs in humans—and there are some indications that it may^{52,53}—the existence of separate initiation and maintenance phases may allow long-term silencing of some targets. Delivery of promoter-targeted siRNA would initiate silencing of a selected disease-causing gene. The RNA-independent maintenance mechanism would then mediate long-term persistence of the silencing effect.

Postscript

The focus of this article should not be taken as special pleading for plants. It is more intended as an attempt to use a set of case histories to justify model systems in general, with plants given an equal footing alongside yeast, worms,

fungi and the rest. I realize that it is not easy for science policy makers and funding agencies to prioritize funding and also that basic scientists—myself included—do not always make their task easy because we often prefer to stay in the comfort zone of basic research. It is easier there than in the uncompromising world of technology, in which the only important consideration is whether something works in the field or patient. Nevertheless, I hope that some of the decision makers in biomedical science will bear in mind the small-RNA story when they decide how to allocate their funds. There will be other discoveries through work in model systems that will have application in diagnosis or treatment of disease.

I hope that this short account may encourage young scientists setting out on their career or deciding which systems to use as they set up their first research group. It may well be that plants will help you address whatever scientific questions you are attracted toward. There is also the possibility, of course, that your efforts with plant systems will help with the small problem of harvesting the sun and feeding the world.

ACKNOWLEDGMENTS

My research has always been a group effort. Without colleagues mentioned here and others, I would not even be considered for this prestigious Lasker Award. I would feel much more comfortable if an award could be made to a research group. Special mention is to be made of A. Hamilton for his commitment to the search for the elusive antisense RNAs. I have been fortunate to receive generous long-term funding from the Gatsby Charitable Foundation that allowed me to respond to the unexpected developments in the early days of my interest in RNA silencing and to follow them over a period of 20 years. I thank the foundation trustees for their confidence and support. R. Freedman alerted Gatsby to my existence, and I will always be grateful to him for that. Other support I have received has come from the UK Biotechnology and Biological Sciences Research Council and from a UK Royal Society Research Professorship. Finally, I acknowledge that I am afflicted with a slightly obsessive streak that must be difficult to live with. I am thankful beyond words to those who have tolerated its consequences.

1. Comfort, N.C. From controlling elements to transposons: Barbara McClintock and the Nobel Prize. *Trends Biochem. Sci.* **26**, 454–457 (2001).
2. Hooke, R. *Micrographia* (Royal Society, London, 1664).
3. van Kammen, A. Beijerinck's contribution to the virus concept — an introduction. *Arch. Virol.* **144**, 1–8 (1999).
4. Pazour, G.J. *et al.* *Chlamydomonas* IFT88 and its mouse homologue, polycystic kidney disease gene *Tg737*, are required for assembly of cilia and flagella. *J. Cell Biol.* **151**, 709–718 (2000).
5. Matzke, M. & Matzke, A.J.M. Planting the seeds of a new paradigm. *PLoS Biol.* **2**, e133 (2004).
6. Zamore, P.D. RNA interference: big applause for silencing in Stockholm. *Cell* **127**, 1083–1086 (2006).
7. Horsch, R.B. *et al.* A simple and general method of

- transferring genes into plants. *Science* **227**, 1229–1231 (1985).
8. Fischhoff, D.A. *et al.* Insect tolerant transgenic tomato plants. *Bio/Technology* **5**, 807–813 (1987).
9. Shah, D.M. *et al.* Engineering herbicide tolerance in transgenic plants. *Science* **233**, 478–481 (1986).
10. Abel, P.P. *et al.* Delay of disease development in transgenic plants that express the tobacco mosaic virus coat protein gene. *Science* **232**, 738–743 (1986).
11. Baulcombe, D.C., Saunders, G.R., Bevan, M.W., Mayo, M.A. & Harrison, B.D. Expression of biologically active viral satellite RNA from the nuclear genome of transformed plants. *Nature* **321**, 446–449 (1986).
12. van Dun, C.M.P., Bol, J.F. & Van Vloten-Doting, L. Expression of alfalfa mosaic virus and tobacco rattle virus coat protein genes in transgenic tobacco plants. *Virology* **159**, 299–305 (1987).
13. Smith, C.J.S. *et al.* Antisense RNA inhibition of polygalacturonase gene expression in transgenic tomatoes. *Nature* **334**, 724–726 (1988).
14. Jorgensen, R. Altered gene expression in plant due to trans interactions between homologous genes. *Trends Biotechnol.* **8**, 340–344 (1990).
15. Napoli, C., Lemieux, C. & Jorgensen, R.A. Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes in *trans*. *Plant Cell* **2**, 279–289 (1990).
16. Smith, C.J.S. *et al.* Expression of a truncated tomato polygalacturonase gene inhibits expression of the endogenous gene in transgenic plants. *Mol. Gen. Genet.* **224**, 477–481 (1990).
17. van der Krol, A.R., Mur, L.A., Beld, M., Mol, J.N.M. & Stuitjij, A.R. Flavonoid genes in petunia: addition of a limited number of gene copies may lead to a suppression of gene expression. *Plant Cell* **2**, 291–299 (1990).
18. Matzke, M.A., Primig, M., Trnovsky, J. & Matzke, A.J.M. Reversible methylation and inactivation of marker genes in sequentially transformed tobacco plants. *EMBO J.* **8**, 643–649 (1989).
19. Longstaff, M., Brigneti, G., Boccard, F., Chapman, S.N. & Baulcombe, D.C. Extreme resistance to potato virus X infection in plants expressing a modified component of the putative viral replicase. *EMBO J.* **12**, 379–386 (1993).
20. English, J.J., Mueller, E. & Baulcombe, D.C. Suppression of virus accumulation in transgenic plants exhibiting silencing of nuclear genes. *Plant Cell* **8**, 179–188 (1996).
21. Lindbo, J.A., Silva-Rosales, L., Proebsting, W.M. & Dougherty, W.G. Induction of a highly specific antiviral state in transgenic plants: implications for regulation of gene expression and virus resistance. *Plant Cell* **5**, 1749–1759 (1993).
22. Potrykus, I. Golden rice and beyond. *Plant Physiol.* **125**, 1157–1161 (2001).
23. Hamilton, A.J., Lycett, G.W. & Grierson, D. Antisense gene that inhibits synthesis of the hormone ethylene in transgenic plants. *Nature* **346**, 284–287 (1990).
24. Hamilton, A.J. & Baulcombe, D.C. A species of small antisense RNA in post-transcriptional gene silencing in plants. *Science* **286**, 950–952 (1999).
25. Hamilton, A., Voinnet, O., Chappell, L. & Baulcombe, D. Two classes of short interfering RNA in RNA silencing. *EMBO J.* **21**, 4671–4679 (2002).
26. Elbashir, S.M. *et al.* Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* **411**, 494–498 (2001).
27. Lee, R.C., Feinbaum, R.L. & Ambros, V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* **75**, 843–854 (1993).
28. Wightman, B., Ha, I. & Ruvkun, G. Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*. *Cell* **75**, 855–862 (1993).
29. Fire, A. *et al.* Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**, 806–811 (1998).
30. Mosher, R.A., Schwach, F., Studholme, D.J. & Baulcombe, D.C. PolIVb influences RNA-directed DNA methylation independently of its role in siRNA

- biogenesis. *Proc. Natl. Acad. Sci. USA* **105**, 3145–3150 (2008).
31. Kasschau, K.D., *et al.* Genome-wide profiling and analysis of *Arabidopsis* siRNAs. *PLoS Biol.* **5**, e57 (2007).
32. Lu, C. *et al.* Elucidation of the small RNA component of the transcriptome. *Science* **309**, 1567–1569 (2005).
33. Vasudevan, S. & Steitz, J.A. AU-rich-element-mediated upregulation of translation by FXR1 and Argonaute 2. *Cell* **128**, 1105–1118 (2007).
34. Covey, S.N., Al-Kaff, N.S., Lángara, A. & Turner, D.S. Plants combat infection by gene silencing. *Nature* **385**, 781–782 (1997).
35. Ratcliff, F., Harrison, B.D. & Baulcombe, D.C. A similarity between viral defense and gene silencing in plants. *Science* **276**, 1558–1560 (1997).
36. Anandalakshmi, R. *et al.* A viral suppressor of gene silencing in plants. *Proc. Natl. Acad. Sci. USA* **95**, 13079–13084 (1998).
37. Brigneti, G. *et al.* Viral pathogenicity determinants are suppressors of transgene silencing in *Nicotiana benthamiana*. *EMBO J.* **17**, 6739–6746 (1998).
38. Kasschau, K.D. & Carrington, J.C. A counterdefensive strategy of plant viruses: suppression of post-transcriptional gene silencing. *Cell* **95**, 461–470 (1998).
39. Voinnet, O., Pinto, Y.M. & Baulcombe, D.C. Suppression of gene silencing: a general strategy used by diverse DNA and RNA viruses. *Proc. Natl. Acad. Sci. USA* **96**, 14147–14152 (1999).
40. Llave, C., Kasschau, K.D. & Carrington, J.C. Virus-encoded suppressor of posttranscriptional gene silencing targets a maintenance step in the silencing pathway. *Proc. Natl. Acad. Sci. USA* **97**, 13401–13406 (2000).
41. Palauqui, J.-C., Elmayan, T., Pollien, J.-M. & Vaucheret, H. Systemic acquired silencing: transgene-specific post-transcriptional silencing is transmitted by grafting from silenced stocks to non-silenced scions. *EMBO J.* **16**, 4738–4745 (1997).
42. Voinnet, O. & Baulcombe, D.C. Systemic signalling in gene silencing. *Nature* **389**, 553 (1997).
43. Wang, X.-H. *et al.* RNA interference directs innate immunity against viruses in adult *Drosophila*. *Science* **312**, 452–454 (2006).
44. Cullen, B.R. Is RNA interference involved in intrinsic antiviral immunity in mammals? *Nat. Immunol.* **7**, 563–567 (2006).
45. Palliser, D. *et al.* An siRNA-based microbicide protects mice from lethal herpes simplex virus 2 infection. *Nature* **439**, 89–94 (2006).
46. Mette, M.F., Aufsatz, W., van der Winden, J., Matzke, M.A. & Matzke, A.J.M. Transcriptional silencing and promoter methylation triggered by double-stranded RNA. *EMBO J.* **19**, 5194–5201 (2000).
47. Volpe, T.A. *et al.* Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. *Science* **297**, 1833–1837 (2002).
48. Pontes, O. *et al.* The *Arabidopsis* chromatin-modifying nuclear siRNA pathway involves a nucleolar RNA processing center. *Cell* **126**, 79–92 (2006).
49. Zhang, X., Henderson, I.R., Lu, C., Green, P.J. & Jacobsen, S.E. Role of RNA polymerase IV in plant small RNA metabolism. *Proc. Natl. Acad. Sci. USA* **104**, 4536–4541 (2007).
50. Herr, A.J., Molnar, A., Jones, A. & Baulcombe, D.C. Defective RNA processing enhances RNA silencing and accelerates flowering in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **103**, 14994–15001 (2006).
51. Jones, L., Ratcliff, F. & Baulcombe, D.C. RNA-directed transcriptional gene silencing in plants can be inherited independently of the RNA trigger and requires Met1 for maintenance. *Curr. Biol.* **11**, 747–757 (2001).
52. Morris, K.V., Chan, S.W.-L., Jacobsen, S.E. & Looney, D.J. Small interfering RNA-induced transcriptional gene silencing in human cells. *Science* **305**, 1289–1292 (2004).
53. Ogawa, Y., Sun, B.K. & Lee, J.T. Intersection of the RNA interference and X-inactivation pathways. *Science* **320**, 1336–1341 (2008).